



Synthesis of Covalent Organic Polymers for removing CO₂ and heavy metal ions with strong affinity

Ko, Dongah; Hwang, Yuhoon; Jakobsen, Mogens Havsteen; Yavuz, Cafer T.; Andersen, Henrik Rasmus

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Ko, D., Hwang, Y., Jakobsen, M. H., Yavuz, C. T., & Andersen, H. R. (2015). *Synthesis of Covalent Organic Polymers for removing CO₂ and heavy metal ions with strong affinity*. Poster session presented at 8th European-Korean Conference on Science and Technology, Strasbourg, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Synthesis of Covalent Organic Polymers for removing CO₂ and heavy metal ions with strong affinity

Dongah Ko¹, Yuhoon Hwang¹, Mogens H. Jakobsen², Cafer T. Yavuz³, Henrik R. Andersen¹


¹ Technical University of Denmark, Department of Environmental Engineering, Miljøvej 113, 2800 Lyngby, ² Technical University of Denmark, Department of Micro- and Nanotechnology, Ørstedss Plads, Bygning 345B, 2800 Kgs. Lyngby, ³ Korea Advanced Institute of Science and Technology (KAIST), Graduate School of EEWs, Daejeon 305-701, Republic of Korea



doko@env.dtu.dk


The demand for functionalized adsorbent that contains not only high surface area but also selectivity and recyclability has increased for several decades. Especially, growing environmental problems such as water pollution and global warming introduced various application possibilities of functionalized adsorbents for pollutant treatment. Our target contaminants are CO₂ and heavy metal ions and they are non-degradable, stable compounds. Hence, adsorption mechanism is considered as a promising solution for removing those pollutants. In this study, we developed several kinds of Covalent Organic Polymers (COPs) and applied them as a functionalized adsorbent for pollutant treatment systems.

1. Covalent Organic Polymers (COPs)



2. Experimental methods

COPs synthesis



Heavy metal ions removal

- Atomic Absorption Spectroscopy
- Sampling time: 10 min, 1 h, 3 h, 24 h, 48 h
- pH measure: Initial point, final point
- BET instrument: volumetric type measurement
- Measurement conditions: ~ 1 bar, 273 K, 298 K
- Selectivity (N₂/CO₂): Ideal Adsorption Solution Theory (IAST) calculation

CO₂ capture

3. Result and Discussion - CO₂ capture

	MEA ^[1]	COP 121		COP 93		
	298K	273K	298K	273K	298K	323K
BET surface area	solution	24.5 m ² g ⁻¹		605.8 m ² g ⁻¹		
CO ₂ uptake (mg g ⁻¹)	60	87.1	59.4	139.6	91.1	60.7
N ₂ uptake (mg g ⁻¹)	NA	5.1	7.5	6.8	4.4	2.3
Selectivity ^[2]	NA	62	29	75.1	73.5	99.2
Thermal stability	110-130°C	up to 450°C		up to 550°C		

[1] Monoethanolamine

[2] Selectivity calculated by IAST calculation

✓ In order to fill up 500 ml bottle with CO₂, 11 g of COP121 or only 7 g of COP 93 are needed.






4. Results and Discussion - Heavy metal ion removal

	COP 6 ^[2]			COP 65 ^[2]		
Cd ²⁺ Initial concentration	0.7 ppm	4 ppm	7 ppm	0.7 ppm	4 ppm	7 ppm
BET surface area	7.300 m ² g ⁻¹			0.001 m ² g ⁻¹		
% removal ^[1] of Cd ²⁺	24%	18%	13%	48%	16%	9%
Langmuir maximum adsorption capacity	1.706 mg/g			0.695 mg/g		
R ²	0.9999			0.9995		

[1] % removal = $\frac{[C]_0 - [C]_e}{[C]_0} \times 100$ C₀ = initial concentration, C_e = equilibrium concentration

[2] Dose amount: 1g/L



5. Summary

